

## 5. References

- ATSDR. 1992. Public health assessment guidance manual. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 1995. Toxicological profile for diethyl phthalate. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 1997. Toxicological profile for di-n-octyl phthalate. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 1998. Toxicological profile for chlorinated dibenzo-p-dioxins: December 1998. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2001a. Toxicological profile for di-n-butyl phthalate. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2001b. Draft guidance manual for preparation of an interaction profile. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2002. Toxicological profile for di(2-ethylhexyl)phthalate. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2004a. Guidance manual for the assessment of joint toxic action of chemical mixtures. Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- ATSDR. 2004b. Interaction profile for benzene, toluene, ethylbenzene, and xylenes (BTEX). Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2004d. Public health assessment guidance manual. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2012. Addendum to the 1998 Toxicological profile for chlorinated dibenzo-p-dioxins. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Agency for Toxic Substances and Disease Registry.
- ATSDR. 2017. Toxicological profile for polybrominated biphenyls and polybrominated diphenyl ethers (update). Atlanta, GA: Agency for Toxic Substances and Disease Registry.
- Aylward LL, Hays SM. 2002. Temporal trends in human TCDD body burden: decreases over three decades and implications for exposure levels. *J Expo Anal Environ Epidemiol* 12:319-328.
- Brown DJ, Van Overneire I, Goeyens L, et al. 2004. Analysis of Ah receptor pathway activation by brominated flame retardants. *Chemosphere* 55(2004):1509-1518.
- CDC. 2006. National Health and Nutrition Examination Survey. Atlanta, GA: CDC/NCEH. Department of Health and Human Services, Centers for Disease Control and Prevention.

- Chen G and Bunce NJ. 2003. Polybrominated diphenyl ethers as Ah receptor agonists and antagonists. *Toxicol Sci* 76:310-320.
- Huang PC, Kuo PL, Guo YL, et al. 2007. Association between urinary phthalate monesters and thyroid hormones in pregnant women. *Hum Reprod* 22(10):2715-2722.
- Koch HM, Presuss R, Angerer J. 2006. Di(2-ethylhexyl)phthalate (DEHP): Human metabolism and internal exposure-- an update and latest results. *Int J Androl* (1):155-165; discussion 181-185.
- Meeker JD, Calafat AM, Hauser R. 2007. Di(2-ethylhexyl) phthalate metabolites may alter thyroid hormone levels in men. *Environ Health Perspect* 115(7):1029-1034.
- Peters AK, van Londen K, Bergman A, et al. 2004. Effects of polybrominated diphenyl ethers on basal and TCDD-induced Ethoxyresorufin activity and cytochrome P450-1A1 expression in MCF-7, HepG2, and H4IIE cells. *Toxicol Sci* 82:488-496.
- Peters AK, Sanderson JT, Bergman A, et al. 2006a. Antagonism of TCDD-induced Ethoxyresorufin-O-deethylation activity by polybrominated diphenyl ethers (PBDEs) in primary cynomolgus monkey (*Macaca fascicularis*) hepatocytes. *Toxicol Lett* 164(2006):123-132.
- Peters AK, Nijmeijer S, Gradin K, et al. 2006b. Interactions of polybrominated diphenyl ethers with the aryl hydrocarbon receptor pathway. *Toxicol Sci* 92(1):133-142.
- Pohl, HR. 2009. Chemical mixtures: Validation of weight-of-evidence predictions for interactions. The 29th International Symposium on Halogenated Persistent Organic Pollutants. August 23-28, 2009, Beijing, China.
- Rider CV, Furr JR, Wilson VS et al. 2010. Cumulative effects of in utero administration of mixtures of reproductive toxicants that disrupt common target tissues via diverse mechanisms of toxicity. *Int J Androl* 33:443-462.
- Sanders JM, Burka LT, Smith CS, et al. 2005. Differential expression of CYP1A1, 2B, and 3A genes in the F344 rat following exposure to a polybrominated diphenyl ether mixture or individual components. *Toxicol Sci* 88(1):127-133.
- Schechter A, Papke O, Chi Tung K, et al. 2005. Polybrominated diphenyl ether flame retardants in the U.S. population: current levels, temporal trends, and comparison with dioxins, dibenzofurans, and polychlorinated biphenyls. *J Occup Environ Med* 47(3):199-211.
- Tanida T, Warita K, Ishihara K, et al. 2009. Fetal and neonatal exposure to three typical environmental chemicals with different mechanisms of action: Mixed exposure to phenol, phthalate, and dioxin cancels the effects of sole exposure on mouse midbrain dopaminergic nuclei. *Toxicol Lett* 189:40-47.
- Tomaszewski, KA., Montgomery CA, Melnick RL. 1988. Modulation of 2,3,7,8-Tetrachlorobenzo-p-dioxin toxicity in F344 rats by di(2-ethylhexyl)phthalate. *Chem Biol Int* 65(1988):205-222.
- Van den Berg M, Birnbaum L, Denison M, et al. 2006. The 2005 World Health Organization reevaluation of human and mammalian toxic equivalency factors for dioxins and dioxin-like compounds. *Toxicol Sci* 93(2):223-241.